

New York State Progress Towards Ending the Epidemic


December 2025

The AIDS Institute uses 16 population-level metrics to help track progress toward Ending the Epidemic. This document shows New York State’s progress to date and goals for each metric. Actual annual outcomes are shown in blue, and goals are shown in purple. The purple dotted line represents the extension of New York State Ending the Epidemic goals to 2024 due to the impact of the COVID-19 pandemic. The COVID-19 pandemic has made an impact on the Ending the Epidemic metrics since 2020.

● Actual ● Goal

1) Estimated New HIV Infections (Incidence)


Significantly reduce the number of new HIV infections



Measure: An estimated number of people who have acquired HIV; diagnosed and undiagnosed (See Note 1 on page 7). Goals are no longer set for incidence due to changes in methodology.
Source: New York State HIV Surveillance System

2) HIV Related Death*

Reduce the percentage of deaths directly related to HIV to 20%




*Metric has delayed reporting to ensure data completeness and accuracy.
Note: Percentages in this metric are based on deaths from all ages.
Measure: The percentage of deaths that were related to HIV among persons with diagnosed HIV. Primary cause of death was used for HIV death ascertainment.
Source: New York State HIV Surveillance System



3) HIV Status Aware


Increase the percentage of persons living with HIV who know their serostatus to at least 98%



Measure: An estimated percentage of people living with HIV who have been diagnosed. (See Note 1 on page 7).
Source: New York State HIV Surveillance System

4) New HIV Diagnoses


Reduce the number of new HIV diagnoses by 55% to 1,515



Note: New diagnoses counts in this metric are based on diagnoses from all ages.
Measure: Number of persons newly diagnosed with HIV.
Source: New York State HIV Surveillance System

5) Newly Diagnosed HIV – Persons with a History of Injection Drug Use

Reduce the percentage of persons newly diagnosed with HIV who indicate a history of injection drug use to 2.8%




Note: Percentages and new diagnoses counts in this metric are based on diagnoses from ages 13 and older.
Measure: Number of persons newly diagnosed with HIV who indicate a history of injection drug use. Includes persons who indicate injection drug use and those who indicate male-to-male sexual contact and injection drug use history.
Source: New York State HIV Surveillance System



Department of Health
AIDS Institute

6) Concurrent AIDS Diagnosis


Reduce the percentage of persons with a diagnosis of AIDS within 30 days of HIV diagnosis to 15%



Note: Percentages and diagnoses counts in this metric are based on diagnoses from ages 13 and older.
Measure: HIV with simultaneous AIDS (stage 3 HIV) diagnosis, or AIDS diagnosis within 30 days of HIV diagnosis.
Source: New York State HIV Surveillance System

7) Linkage to Care After Diagnosis


Increase the percentage of newly diagnosed persons linked to HIV medical care within 30 days of diagnosis to at least 90%



Note: Percentages in this metric are based on diagnoses from ages 13 and older.
Measure: Newly diagnosed with any viral load, CD4, or genotype test within 30 days of diagnosis by diagnosis year.
Source: New York State HIV Surveillance System

8) Receiving HIV Medical Care

Increase the percentage of persons living with diagnosed HIV who receive HIV medical care to 90%




Note: Percentages in this metric are based on individuals living with diagnosed HIV aged 13 and older.
Measure: Any viral load, CD4, or genotype test in a calendar year.
Source: New York State HIV Surveillance System



Department of Health
AIDS Institute

9) Viral Load Suppression – Newly Diagnosed HIV

Increase the percentage of persons newly diagnosed with HIV who reach viral load suppression within 3 months of diagnosis to 75%




Note: Percentages in this metric are based on diagnoses from ages 13 and older.

Measure: Viral load test suppressed (non-detectable or <200 copies per milliliter) within 91 days from the date of HIV diagnosis.

Source: New York State HIV Surveillance System

10) Viral Load Suppression – Persons Living with Diagnosed HIV

Increase the percentage of persons living with diagnosed HIV with suppressed viral load to 85%




Note: Percentages in this metric are based on individuals living with diagnosed HIV aged 13 and older.

Measure: Last viral load test in calendar year is non-detectable or <200 copies per milliliter.

Source: New York State HIV Surveillance System

11) Viral Load Suppression – Receiving HIV Medical Care

Increase the percentage of people living with diagnosed HIV who receive HIV medical care with suppressed viral load to 95%



Note: Percentages in this metric are based on individuals living with diagnosed HIV aged 13 and older.

Measure: Last viral load test in calendar year is non-detectable or <200 copies per milliliter, among those in care during the calendar year.


Source: New York State HIV Surveillance System



Department of Health
AIDS Institute

12) Sustained Viral Load Suppression

Increase the percentage of persons living with diagnosed HIV with sustained viral suppression to 75%




Note: Percentages in this metric are based on individuals living with diagnosed HIV aged 13 and older.

Measure: Viral load test suppressed (non-detectable or <200 copies per milliliter) on all viral load tests in the previous 2 years, among those with at least 2 viral load tests in the previous 2 years.

Source: New York State HIV Surveillance System

13) Time to AIDS Diagnosis*

Reduce the rate at which persons diagnosed with HIV progress to AIDS by 50%



*Metric has delayed reporting to ensure data completeness and accuracy


Note: Percentages in this metric are based on diagnoses from ages 13 and older.

Measure: AIDS diagnosis within 2 years of HIV diagnosis.

Source: New York State HIV Surveillance System

14) Stigma*

Decrease stigma experienced among persons living with diagnosed HIV to at least 25



*Metric has delayed reporting to ensure data completeness and accuracy

Measure: The weighted median score on a 10-item scale ranging from 0 (no stigma) to 100 (high stigma) that measures 4 dimensions of HIV stigma (personalized stigma during the past 12 months, disclosure concerns, negative self-image, and perceived public attitudes about persons with HIV).


Source: New York State Medical Monitoring Project



Department of Health
AIDS Institute

15) Pre-Exposure Prophylaxis (PrEP) Utilization


Increase the number of individuals filling prescriptions for Pre-Exposure Prophylaxis (PrEP) to 65,000



Measure: Number of individuals filling at least one prescription for Truvada or Descovy within the calendar year.
Source: IDV® (Integrated Dataverse) from Symphony Health

16) Pre-Exposure Prophylaxis (PrEP) Utilization – Medicaid

Increase the number of Medicaid recipients filling prescriptions for Pre-Exposure Prophylaxis (PrEP) to 30,000



Measure: Number of Medicaid recipients filling at least one prescription for Truvada or Descovy within the calendar year.
Source: IDV® (Integrated Dataverse) from Symphony Health & New York State Medicaid Data Warehouse



Ending the Epidemic Metrics ● Actual ● Goal ● Reached goal

	1 Estimated New HIV Infections	2 HIV Related Death*	3 HIV Status Aware	4 New HIV Diagnoses	5 Newly Diagnosed HIV– Persons with a History of Injection Drug Use	6 Concurrent AIDS Diagnosis	7 Linkage to Care after Diagnosis	8 Receiving HIV Medical Care
	#	%	%	#	%	#	%	%
2012								
2013	2,895	36%	91%	3,391	4.8%	181	21.7%	736
2014	2,803	32%	91%	3,443	4.4%	158	19.6%	674
2015	2,587	34%	92%	3,155	4.2%	150	19.4%	612
2016	2,426	31%	92%	2,881	3.9%	127	18.8%	541
2017	2,201	27%	92%	2,769	4.0%	106	19.8%	547
2018	1,965	26%	93%	2,481	4.9%	122	20.9%	519
2019	1,846	24%	93%	2,377	4.5%	108	19.6%	465
2020	1,764	18%	93%	1,933	3.3%	63	19.1%	369
2021	1,517	18%	93%	2,123	4.1%	86	20.5%	436
2022	1,681	18%	94%	2,318	3.1%	72	18.0%	418
2023	1,615	18%	94%	2,517	4.9%	124	19.4%	488
2024	1,986	20%	93%	2,555	4.3%	109	20.7%	529

	9 Viral Load Suppression– Newly Diagnosed HIV	10 Viral Load Suppression – Persons Living with Diagnosed HIV	11 Viral Load Suppression– Receiving HIV Medical Care	12 Sustained Viral Load Suppression	13 Time to AIDS Diagnosis*	14 Stigma*	15 Pre-Exposure Prophylaxis (PrEP) Utilization	16 Pre-Exposure Prophylaxis (PrEP) Utilization – Medicaid
	%	%	%	%	%	Score	#	#
2012					10.4%			
2013	23%	66%	81%	54%	6.8%			251
2014	29%	68%	84%	58%	7.1%		3,388	650
2015	32%	67%	85%	62%	6.9%	34	9,324	2,244
2016	39%	70%	87%	64%	5.9%	35	17,368	4,226
2017	48%	72%	87%	65%	5.0%	38	24,284	6,026
2018	55%	75%	89%	67%	6.6%	30	31,964	8,085
2019	59%	77%	89%	69%	5.7%	29	39,921	9,447
2020	54%	75%	87%	68%	4.1%	28	39,282	9,460
2021	54%	78%	89%	70%	4.4%	29	44,046	11,671
2022	58%	79%	90%	71%	5.0%	31	50,558	12,428
2023	58%	80%	90%	72%	5.1%	28	55,891	13,514
2024	61%	81%	91%	72%	5.1%	25	60,313	13,480

*Metric has delayed reporting to ensure data completeness and accuracy

Note 1: Incidence and undiagnosed estimates are calculated using Centers for Disease Control and Prevention methodology released in 2024.

For more information, please contact the
Data Analysis and Research Translation team
at BHA@health.ny.gov



Department of Health
AIDS Institute